



Europäisches Patentamt
European Patent Office
Office européen des brevets

Publication number:

**0 293 530
A2**

(12)

EUROPEAN PATENT APPLICATION

(21) Application number: **87308721.7**

(51) Int. Cl. 4: **C07K 7/20 , A61K 37/02 ,
C12P 21/00 , C12N 15/00**

(22) Date of filing: **01.10.87**

(30) Priority: **05.06.87 GB 8713240**

(43) Date of publication of application:
07.12.88 Bulletin 88/49

(84) Designated Contracting States:
AT BE CH DE ES FR GB GR IT LI LU NL SE

(71) Applicant: **Proteus Biotechnology Limited
41 Park Lane Poynton
Stockport Cheshire SK12 1RD(GB)**

(72) Inventor: **Morrison, Christopher
Route de Belfeux 28 Chin 1711
Corminboeuf, F.R.(CH)
Inventor: Robson, Barry
26 Jessop Drive Marple
Stockport Cheshire(GB)
Inventor: Fishleigh, Robert
14 Clifton Road
Chorlton Manchester(GB)**

(74) Representative: **Hildyard, Edward Martin et al
Frank B. Dehn & Co. Imperial House 15-19
Kingsway
London WC2B 6UZ(GB)**

(54) Improvements in or relating to hormones.

(57) An analogue of lutenising hormone-releasing hormone (LHRH) having the amino acid sequence:
pGlg-His-Trp-Ser-Tyr-X-Len-Arg-Pro-Gly-Y-Z

wherein X represents Gly or a D-amino acid, Y represents one or more amino acid residues, which may be the same or different, and Z represents Cys or Tyr, such that the solution conformation of said analogue is substantially similar to that of native LHRH.

The analogues may be used to prepare vaccines for reducing fertility in mammals.

EP 0 293 530 A2

Improvements in or relating to hormones

The present invention relates to extended analogues of the mammalian hormone referred to as luteinising hormone-releasing hormone (LHRH). In particular, it relates to analogues of LHRH formed by adding to the C-terminus of the native LHRH amino acid sequence a short additional amino acid sequence ending with a cysteine residue, such that the conformation in solution is substantially unchanged, and polypeptide conjugates thereof suitable for raising anti-LHRH antibodies.

LHRH is a decapeptide having the amino acid sequence

pGlg-His-Trp-Ser-Tyr-Gly-Leu-Arg-Pro-GlyNH₂ wherein pGlg represents a pyroglutamate residue and GlyNH₂ represents glycine-amide. In mammals, this hormone is secreted by the hypothalamus and is responsible for controlling the release of luteinising hormone and follicle-stimulating hormone from the pituitary (Schally et al., *Science* (1973) 179, 341-350; Guillemin, R., *Contraception* (1972) 5, 1-32). The latter hormones are directly involved in controlling the development and the normal physiological functioning of testes and ovaries.

Active immunisation of various mammals to LHRH, eg, rats, rabbits, sheep and cattle, has been shown to lead in the case of males to testicular regression, reduction of testosterone secretion and cessation of spermatogenesis and in the case of females to loss of cycling and ovarian regression, and it has long been appreciated that active immunisation to LHRH could provide a desirable means for routinely reducing fertility in domestic mammals and farm animals, particularly bull calves. In a comparative study of steer calves and immunologically-castrated calves produced by active immunisation to LHRH, the so-called "immunocastrates" were found to be superior in terms of their carcass composition since in these animals anabolic influence of the testes was not totally abolished (Robertson et al., *Veterinary Record* (1982) 111, 529-531). Moreover, surgical castration involves a degree of risk of infection and trauma, particularly if attempted on other than very young animals. Nevertheless, active immunisation to LHRH using a polypeptide conjugate of LHRH has previously required deleterious adjuvants and moreover the polypeptide conjugates themselves have been immunogenically unsatisfactory.

Hitherto published procedures for the conjugation of LHRH to a polypeptide carrier, e.g. bovine or human serum albumin, tetanus toxoid or thyroglobulin, have generally involved condensation of the hormone with a water-soluble carbodiimide. The absence of free carboxyl or amino moieties in

LHRH means that coupling in such methods is probably effected through the hydroxyl group of Ser⁴ or via a carboxymethylated derivative of His². As a result, the immunogen is poorly-defined and unlikely to retain all the structural features of free LHRH in solution as considered desirable from the point of view of obtaining anti-LHRH antibodies capable of blocking functioning of LHRH *in vivo*; there is a danger that the peptide is attached through a region important for immunological recognition. Effective immunisation of mammals using such conjugates to provide a high titre of anti-LHRH antibodies capable of significantly reducing the biological efficacy of endogenous LHRH has only been achieved in the presence of an adjuvant liable to cause undesirable side effects, most commonly Freund's complete or incomplete adjuvant. Freund's complete adjuvant interferes with the tuberculin test in cattle and in addition this adjuvant and also Freund's incomplete adjuvant cause a variable amount of chronic inflammatory reaction at the site of injection. Such procedures have accordingly achieved no practical importance.

A totally synthetic LHRH vaccine based on muramyl dipeptide (MDP) has recently been shown to be capable of effecting immunological castration in male mice (Carelli et al., *Proc. Nat. Acad. Sci. USA* (1982) 79, 5392-5395; Carelli et al., *Int. J. Immunopharmacol.* (1985) 7, 215-224). However, due to the apparent pyrogenic effects of MDP, this technique of immunological castration is also likely to prove unacceptable for general use in veterinary practice.

With the aim of overcoming the above-mentioned problems, we have now designed analogues of LHRH with a short peptide extension at the C-terminus of the native amino acid sequence, which we have predicted by potential energy calculations to have substantially the same conformation as native LHRH in solution and may be readily linked to a polypeptide carrier via the side chain of a cysteine or tyrosine residue provided at the C-terminus.

According to one aspect of the present invention, we thus provide an analogue of LHRH having the amino acid sequence

pGlg-His-Trp-Ser-Tyr-X-Leu-Arg-Pro-Gly-Y-Z wherein X represents Gly or a D-amino acid, Y represents one or more amino acid residues, which may be the same or different, and Z represents Cys or Tyr, such that the solution conformation of said analogue is substantially similar to that of native LHRH.

Preferably, the C-terminal extension consists of 1-3 glycine residues followed by a cysteine or

tyrosine residue at the C-terminus, most preferably a single glycine residue followed by a cysteine residue (the analogue of this type wherein the native LHRH sequence is joined to a Gly-Cys extension will hereinafter be referred to as LHRH-Gly-Cys-OH). Computer-aided potential energy calculations have indicated that these preferred LHRH analogues preferably adopt a conformation in solution virtually identical to the type II' turn around Gly⁶-Leu⁷ predicted as the lowest energy solution conformation for native LHRH.

By substituting for the Gly residue at position 6 in an extended LHRH analogue as above a D-amino acid, e.g. D-Ala, D-Lys or D-Ser(tBu), further advantageous extended LHRH analogues may be obtained without radical alteration of the solution conformation. Such analogues may be expected to have the benefit of improved stability to degradation *in vivo*.

The novel analogues may be synthesised by standard peptide synthesis techniques, e.g. by the Fmoc-polyamide method using the solid phase resin developed by Arshady, R. et al. (J. Chem. Soc. Perkin Trans. (1981) 1, 529-537) and fluorenylmethoxycarbonyl (Fmoc) protection of the individual amino acids incorporated (Atherton, E. et al., J. Chem. Soc. Perkin Trans. (1983) 1, 65-73).

The novel analogues may also be synthesised indirectly by DNA synthesis and conventional techniques of genetic engineering.

Another aspect of the present invention therefore provides DNA molecules coding for at least one extended LHRH analogue as described herein preferably incorporated into a suitable expression vector replicable in microorganisms or in mammalian cells.

According to a further aspect of the present invention, we provide an extended LHRH analogue as hereinbefore described conjugated to a polypeptide carrier e.g. via the side chain of the C-terminal cysteine or tyrosine residue. Formation of such an LHRH analogue-carrier conjugate may conveniently be achieved by modification of one or more lysine side chains of the carrier polypeptide with maleimide, e.g. by treatment with N-γ-maleimidobutyryloxysuccinimide (Calbiochem), and subsequent coupling of the LHRH analogue via the side chain of the C-terminal cysteine residue to give a thioether bond. Analogues having a terminal tyrosine residue may be coupled to a carrier polypeptide e.g. by a diazo bond formed with N-(4-diazophenyl) maleimide. Other coupling reactions and reagents may of course be used to equivalent effect. The conjugate may also, in principle, be produced via a DNA molecule coding therefor which is preferably incorporated into a suitable expression vector replicable in microorganisms or in mammalian cells.

We believe that in such conjugates the LHRH moiety substantially retains the free solution conformation which is optimal for production of anti-LHRH antibodies, since the linkage is both flexible and of sufficient length to hold the peptide away from the surface of the carrier.

By employing conjugates of LHRH analogues of this type, it has been found possible to achieve effective adjuvant-free immunisation to LHRH in mammals and we consider this technique to be a highly desirable alternative to surgical castration for veterinary control of reproductive function, e.g. in dogs, cats and horses. Moreover, in the case of humans having an androgen-sensitive or oestrogen-sensitive carcinoma, e.g. prostate carcinoma or mammary carcinoma, use of such conjugates to induce anti-LHRH antibody production and thus lower steroid levels may also be beneficial.

For immunisation of a wide range of mammals to LHRH, examples of preferred carriers include purified protein derivative of tuberculin (Central Veterinary Laboratory, Weybridge, U.K.) and tetanus toxoid (Wellcome), purified protein derivative of tuberculin being most favoured. When using purified protein derivative of tuberculin (commonly referred to as PPD) as the carrier for production of anti-LHRH antibodies, in order to achieve a high titre of the required antibodies it is desirable for the recipient of the LHRH analogue - PPD conjugate to be tuberculin sensitive, e.g. by virtue of an earlier BCG vaccination, as previously described for antibody production to other hapten-PPD conjugates (Lachmann et al. in "Synthetic Peptides as Antigens", Ciba Foundation Symposium 119 (1986) pp. 25-40). In the U.K. and many other countries, the population is offered BCG vaccination and is therefore largely PPD-sensitive. Hence, LHRH analogue-PPD conjugates of the present invention are considered highly advantageous for use in the clinical control of carcinoma in humans.

According to a still further aspect of the present invention, we provide a method of reducing fertility in a mammal which comprises immunising said mammal with an extended LHRH analogue or with a carrier conjugate thereof as hereinbefore described so as to produce a titre of anti-LHRH antibodies sufficient to significantly reduce the biological efficacy of endogenous LHRH. Normally no adjuvant is necessary. In a particularly preferred embodiment of this method, an LHRH analogue-PPD conjugate of the present invention is administered intradermally in isotonic saline, if necessary after priming of the recipient with BCG vaccine to achieve tuberculin sensitivity. For such an immunisation protocol, it is particularly desirable to employ the thioether conjugate of maleimide - modified PPD and LHRH-Gly-Cys-OH (hereinafter

referred to as LHRH-Gly-Cys-OH-PPD).

The following non-limiting example further illustrates the present invention.

Example

1. Solution conformation analysis of LHRH-Gly-Cys-OH

The conformational preferences in solution of LHRH and LHRH-Gly-Cys-OH (see Figures 1(a) and 1(b)) were studied using the LUCIFER program for potential energy minimisation (Robson and Platt, J. Mol. Biol. (1986) 188, 259-281). The program was run on a CDC Cyber 205 computer and solvent effects were modelled using the representation previously used for a study of neurotensin (Ward et al., Regul. Peptides (1986) 15, 197).

From energy-minimisation studies, we have discovered that LHRH-Gly-Gly-Cys-OH and LHRH-Gly-Gly-Gly-Cys-OH also have very similar solution conformations to LHRH.

2. Preparation of LHRH-Gly-Cys-OH-PPD and other LHRH analogue-carrier conjugates.

Bovine PPD (10mg, Batch no. 291, Central Veterinary Laboratory, Weybridge, UK) was dissolved in 0.5ml of 0.05M NaKPO₄, 0.14M NaCl, pH 7.0 (buffer A) and treated with 1mg N-γ-maleimidobutyryloxysuccinimide (Calbiochem) previously dissolved in 5ul freshly distilled, dry dimethyl formamide. The mixture was stirred for 1 hour at 23°C and applied to a column of Sephadex G25 (0.9 x 25cm) equilibrated with buffer A. The modified PPD eluting in the excluded volume was transferred to a stoppered vessel and to it was added dropwise, with stirring, 10mg of LHRH-Gly-Cys-OH (85% purity) dissolved in 1ml of buffer A previously degassed and purged with nitrogen. The mixture was stirred at 23°C under nitrogen for 2 hours and the amount of LHRH which bound to the PPD determined by estimating the free thiol content of the mixture at regular intervals using 5,5'-dithiobis(2-nitrobenzoic acid) (Sigma) according to the method of Deakin et al. (Biochem. J. (1963) 89, 296-304). A coupling efficiency of 65% was noted. The conjugate was extensively dialysed against several changes of distilled water at 4°C, lyophilised, and stored at -20°C before use.

Tetanus toxoid (TT) (Wellcome) and bovine serum albumin (BSA) (Sigma) were conjugated to LHRH-Gly-Cys-OH exactly as described for PPD above.

3. Immunisation studies with rats

(a) Immunisation schedule

Groups of 5 male rats (AO x DA, F₁ rats, 4 months old) were immunised with an amount of conjugate equivalent to 50ug LHRH-Gly-Cys-OH according to the following schedule. Groups 1 and 2 received LHRH-Gly-Cys-OH-PPD intradermally in saline. Groups 3 and 4 received LHRH-Gly-Cys-OH-PPD as an alum precipitate subcutaneously. Groups 5 and 6 received LHRH-Gly-Cys-OH-TT subcutaneously in alum or Freund's complete adjuvant (FCA) respectively. Group 7 received no immunisation. Groups 1 and 3 received BCG vaccine (Glaxo) equivalent to one half the human dose one month prior to the first immunisation with conjugate. Immunisation consisted of a primary injection followed by two booster injections at one monthly intervals.

The animals in the study were subsequently bled by cannulation of the tail artery at fortnightly intervals for estimation of antibody titres, and then were sacrificed 3 months after the primary immunisation, the testes removed, weighed and subjected to histological examination.

Estimation of anti-LHRH antibody titre was achieved by ELISA using plastic microtitre plates coated with LHRH-Gly-Cys-OH-BSA. Assays were developed with 1:1000 dilution of goat anti-rat 1gG alkaline phosphatase conjugate.

Removed testes were fixed in neutral buffered formalin, embedded in paraffin wax and 5 micron sections cut. Sections were stained with haematoxylin and eosin.

(b) Results

(i) Antibody titres to LHRH

Each of the groups of rats immunised with LHRH-Gly-Cys-OH conjugates gave an antibody response to the hormone detectable by ELISA. Figure 2 shows the antibody titres at week 10 for each group. Immunisation with the conjugate of LHRH-Gly-Cys-OH-TT led to significant antibody levels, and these were higher in the group receiving conjugate in FCA. Although all four groups of rats receiving LHRH-Gly-Cys-OH-PPD gave antibody responses, the responses in those groups (1 and 3) previously primed with BCG, were considerably higher than in those which were not (2 and 4). LHRH-Gly-Cys-OH-PPD given intradermally was a

very poor immunogen in normal rats, but in the BCG primed animals led to the highest titre of antibody seen in any of the groups. LHRH-Gly-Cys-OH-PPD absorbed onto alum and given subcutaneously led to significant antibody production in normal rats, but in the BCG primed rats the response was inferior to that obtained with the soluble antigen given intradermally.

(ii) Physiological effects of immunisation to LHRH

Mean testis weights for individual rats in each group are shown in Figure 3. Marked testicular regression was apparent in all members of the group receiving soluble LHRH-Gly-Cys-OH-PPD after BCG priming and also in the group treated with LHRH-Gly-Cys-OH-TT in FCA although, here, one animal showed no effects. Less marked effects were seen in Group 3 (2 of 5) and Group 5 (1 of 5). Histological examination of testes revealed that in those individuals exhibiting severe regression there was evidence of marked atrophy of seminiferous tubules and absence of spermatogenesis.

In further studies, a conjugate of LHRH and PPD prepared using a water soluble carbodiimide in conventional manner was shown to be unable to induce a physiological response when used to immunise rats subcutaneously or intradermally in the absence of an adjuvant. D-Lys⁶-LHRH conjugated to PPD with glutaraldehyde was likewise found to be ineffective.

Claims

1. An analogue of lutenising hormone-releasing hormone (LHRH) having the amino acid sequence:

pGlg-His-Trp-Ser-Tyr-X-Len-Arg-Pro-Gly-Y-Z wherein X represents Gly or a D-amino acid, Y represents one or more amino acid residues, which may be the same or different, and Z represents Cys or Tyr, such that the solution conformation of said analogue is substantially similar to that of native LHRH.

2. An analogue as claimed in claim 1 wherein X represents Gly.

3. An analogue as claimed in claim 1 wherein X represent D-Ala, D-Lys or D-Ser(tBu).

4. An analogue as claimed in any of claims 1-3 wherein Y represents 1-3 glycine residues.

5. An analogue as claimed in claim 4 wherein Y represents one glycine residue and Z represents Cys.

6. An analogue as claimed in any of claims 1-5 conjugated to a polypeptide vaccine carrier.

7. An analogue-carrier conjugate as claimed in claim 6 wherein said carrier is purified protein derivative of tuberculin or tetanus toxoid.

8. An analogue-carrier conjugate as claimed in claim 6 or 7 wherein said conjugation is via the side-chains of the C-terminal cysteine residue.

9. A vaccine comprising an LHRH analogue or analogue-carrier conjugate as claimed in any of claims 1 to 8 and a pharmaceutically acceptable excipient.

10. A DNA molecule coding for an analogue or analogue-carrier conjugate as claimed in any of claims 1 to 8.

11. A DNA molecule as claimed in claim 10 incorporated into an expression vector replicable in microorganisms or in mammalian cells.

12. A method of reducing fertility in a mammal which comprises immunising said mammal with an LHRH analogue or analogue-carrier conjugate as claimed in any of claims 1 to 8 so as to produce a titre of anti-LHRH antibodies sufficient to reduce significantly the biological efficacy of endogenous LHRH.

13. The use of an LHRH analogue or analogue-carrier conjugate as claimed in any of claims 1 to 8 for the preparation of a composition for reducing fertility in a mammal.

14. The use of an LHRH analogue or analogue-carrier conjugate as claimed in any of claims 1 to 8 for the preparation of a composition for the clinical control of androgen-sensitive or oestrogen-sensitive carcinoma.

15. A process for the preparation of an LHRH analogue or analogue-carrier conjugate as claimed in any of claims 1 to 8, which comprises:

(a) synthesising said LHRH analogue by a peptide synthesis technique known per se; or


(b) cultivating microorganisms or mammalian cells containing an expression vector incorporating DNA coding for said LHRH analogue;

whereafter if desired the analogue is conjugated to a polypeptide vaccine carrier by methods known per se.

FIG. 1.

Stereoscopic diagrams of the predicted lowest energy solution conformers of (a) mammalian LHRH, and (b) the LHRH-Gly-Cys-OH analogue.

a



b




FIG. 2.

Day 70 anti-LHRH-Gly-Cys-OH antibody titres in rats immunised with various LHRH-Gly-Cys-OH conjugates.





FIG. 3.

Effect of immunisation with various LHRH-Gly-Cys-OH conjugates on testicular weight in rats.



19



Europäisches Patentamt
European Patent Office
Office européen des brevets

11 Publication number:

**0 293 530
A3**

12

EUROPEAN PATENT APPLICATION

21 Application number: **87308721.7**

51 Int. Cl.⁵: **C07K 7/20, A61K 37/02,
C12P 21/00, C12N 15/00**

22 Date of filing: **01.10.87**

30 Priority: **05.06.87 GB 8713240**

43 Date of publication of application:
07.12.88 Bulletin 88/49

84 Designated Contracting States:
AT BE CH DE ES FR GB GR IT LI LU NL SE

88 Date of deferred publication of the search report:
30.05.90 Bulletin 90/22

71 Applicant: **Proteus Biotechnology Limited**
41 Park Lane Poynton
Stockport Cheshire SK12 1RD(GB)

72 Inventor: **Morrison, Christopher**
Route de Belfeux 28 Chin 1711
Corminboeuf, F.R.(CH)
Inventor: **Robson, Barry**
26 Jessop Drive Marple
Stockport Cheshire(GB)
Inventor: **Fishleigh, Robert**
14 Clifton Road
Chorlton Manchester(GB)

74 Representative: **Hildyard, Edward Martin et al**
Frank B. Dehn & Co. Imperial House 15-19
Kingsway
London WC2B 6UZ(GB)

54 Improvements in or relating to hormones.

57 An analogue of lutenising hormone-releasing hormone (LHRH) having the amino acid sequence:
pGlg-His-Trp-Ser-Tyr-X-Len-Arg-Pro-Gly-Y-Z

wherein X represents Gly or a D-amino acid, Y represents one or more amino acid residues, which may be the same or different, and Z represents Cys or Tyr, such that the solution conformation of said analogue is substantially similar to that of native LHRH.

The analogues may be used to prepare vaccines for reducing fertility in mammals.

EP 0 293 530 A3



European Patent
Office

PARTIAL EUROPEAN SEARCH REPORT

which under Rule 45 of the European Patent Convention shall be considered, for the purposes of subsequent proceedings, as the European search report

Application number

EP 87 30 8721

DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int. Cl. 4)
X	FEBS LETTERS, vol. 214, no. 1, 6th April 1987, pages 65-70, Elsevier Science Publishers B.V., Amsterdam, NL; C.A. MORRISON et al.: "Computer-aided design and physiological testing of a luteinising hormone-releasing hormone analogue for "adjuvant-free" immunocastration"	1-11, 13-15	C 07 K 7/20 A 61 K 37/02 C 12 P 21/00 C 12 N 15/00
A	* Whole article * -- CHEMICAL ABSTRACTS, vol. 106, no. 17, 27th April 1987, abstract no. 134121f, page 298, Columbus, Ohio, US; T.E. PALEN et al.: "Characterization of an enzyme that is capable of processing pro-gonadotropin-releasing hormone protein", & ARCH. BIOCHEM. BIOPHYS. 1986, 251(2), 543-50 ./.		

TECHNICAL FIELDS SEARCHED (Int. Cl. 4)

A 61 K
C 07 K
C 12 N
C 12 P

INCOMPLETE SEARCH

The Search Division considers that the present European patent application does not comply with the provisions of the European Patent Convention to such an extent that it is not possible to carry out a meaningful search into the state of the art on the basis of some of the claims.

Claims searched completely: 1-11, 13-15

Claims searched incompletely:

Claims not searched: 12
Reason for the limitation of the search:

Method for treatment of the human or animal body by surgery or therapy (See art. 52(4) of the European Patent Convention).

Place of search The Hague	Date of completion of the search 01-02-1990	Examiner MASTURZO
CATEGORY OF CITED DOCUMENTS X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document		

EPO Form 1505.1 03/82



DOCUMENTS CONSIDERED TO BE RELEVANT			CLASSIFICATION OF THE APPLICATION (Int. Cl. 4)
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	
A	<p>* Abstract *</p> <p>--</p> <p>CHEMICAL ABSTRACTS, vol. 106, no. 21, 25th May 1987, page 99, abstract no. 169348n, Columbus, Ohio, US;</p> <p>T.J. GORCS et al.: "Possible recognition of the GnRH receptor by an antiserum against a peptide encoded by nucleotide sequence complementary to mRNA of a GnRH precursor peptide", & PEPTIDES (FAYETTEVILLE, N.Y.) 1986, 7(6), 1137-45</p>	1	
	<p>* Abstract *</p> <p>--</p> <p>FR-A-2 465 486 (ROUSSEL-UCLAF)</p> <p>* Whole document, especially the description on pages 1 and 2 *</p> <p>-----</p>	1 1-3, 13-14	TECHNICAL FIELDS SEARCHED (Int. Cl. 4)

THIS PAGE BLANK (USPTO)